



# Phytoremediation of Cr(VI) polluted soil

**Marius Gheju**

**“Politehnica” University of Timisoara, Faculty of Industrial Chemistry and Environmental Engineering, Bd. V. Parvan Nr. 6, Et. 4, Timisoara, 300223, Romania**

**e-mail: [marius.gheju@chim.upt.ro](mailto:marius.gheju@chim.upt.ro)**

**Ionel Balcu**

**National Institute for Development in Electrochemistry and Condensed Matter,  
Str. Dr. Aurel Păunescu Podeanu Nr. 144, Timisoara, 300587, Romania**

**e-mail: [ionel\\_balcu@yahoo.com](mailto:ionel_balcu@yahoo.com)**

# Chromium natural occurrence

	Abundance
<b>Universe</b>	<b>15,000 ppb</b>
<b>Sun</b>	<b>20,000 ppb</b>
<b>Meteorites</b>	<b>3,000 ppm</b>
<b>Air</b>	<b>0.01-0.03 <math>\mu\text{g}/\text{m}^3</math></b>
<b>Sea water</b>	<b>0.3 ppb</b>
<b>Rivers</b>	<b>1 ppb</b>
<b>Continental crust</b>	<b>80-200 ppm</b>
<b>Ultramafic igneous rocks</b>	<b>1,000-3,400 ppm</b>
<b>Basaltic igneous rocks</b>	<b>40-600 ppm</b>
<b>Granitic igneous rocks</b>	<b>2-90 ppm</b>
<b>Shales and clays</b>	<b>30-590 ppm</b>
<b>Coals</b>	<b>10-1,000 ppm</b>

# Chromium natural occurrence in food

<b>Food</b>	<b>Mean concentration (<math>\mu\text{g}/\text{kg}</math>)</b>
<b>Fresh vegetables</b>	<b>30-140</b>
<b>Fresh fruits</b>	<b>90-190</b>
<b>Dairy products</b>	<b>100</b>
<b>Chicken eggs</b>	<b>160-520</b>
<b>Whole fish</b>	<b>50-80</b>
<b>Meat</b>	<b>110-230</b>
<b>Seafoods</b>	<b>120-470</b>
<b>Grains and cereals</b>	<b>40-220</b>
<b>Refined sugar</b>	<b>20</b>
<b>Cigarette tobacco</b>	<b>240-14,600</b>

# Chromium natural occurrence in human body

<b>Tissue / fluid</b>	<b>Mean content</b>
<b>Human body</b>	<b>30 µg/kg (dry weight)</b>
<b>Serum</b>	<b>0.006 µg/L</b>
<b>Urine</b>	<b>0.4 µg/L</b>
<b>Breast milk</b>	<b>0.3 µg/L</b>
<b>Lung</b>	<b>203 µg/kg (wet weight)</b>
<b>Hair</b>	<b>0.234 mg/kg</b>
<b>Nail</b>	<b>0.52 mg/kg</b>

# Sources of chromium soil contamination

- Chrome electroplating;
- Wood preserving;
- Leather tanning;
- Manufacturing of dyes and pigments;
- Textile industries (dyes);
- Disposal of commercial products containing chromium;
- Disposal of chromium waste from industry;
- Disposal of coal ash from electric utilities;
- Combustion processes (burning of coal and oil);
- Sewage sludge and municipal refuse incineration;
- Steel production;
- Chemical manufacturing and use of compounds containing chromium;
- Refractory production;
- Cement production.

# Chromium in the environment

Although chromium can exist in oxidation states ranging from (-IV) to (+VI), in natural environments Cr exists in two main oxidation states: Cr(+VI) and Cr(+III), characterized by different chemical behaviour and toxicity.

# Chromium (VI) chemical behaviour

pH < 6.5:  $\text{HCrO}_4^- / \text{Cr}_2\text{O}_7^{2-}$


pH > 6.5:  $\text{CrO}_4^{2-}$

pH < 0:  $\text{H}_2\text{CrO}_4, \text{Cr}_3\text{O}_{10}^{2-}$


$\text{Cr}_4\text{O}_{13}^{2-}$

**Significant solubility**

**Significant mobility**



Distribution of Cr(VI) species as function of pH  
 $C_{\text{Cr(VI)}} = 5 \text{ mmol/L}$




Distribution of Cr(VI) species as function of total Cr(VI) concentration, pH 4

# Chromium (III) chemical behaviour

pH < 3.0:  $\text{Cr}^{3+}$   
pH = 3.0 - 5.5:  $\text{Cr}(\text{OH})^{2+}$   
pH = 5.5 - 6.0:  $\text{Cr}(\text{OH})_2^+$   
pH = 6.0 - 11.5:  $\text{Cr}(\text{OH})_3, \text{Cr}_2\text{O}_3$   
pH > 11.5:  $\text{Cr}(\text{OH})_4^-$

**Low solubility**

**Low mobility**



# Chromium (VI) toxicological behaviour

- Highly toxic to humans, animals, plants and microorganisms;
- Known human carcinogen by the inhalation route of exposure;
- Animal carcinogen by the oral route of exposure.

# Chromium (III) toxicological behaviour

- Less toxic than Cr(VI);
- No evidence of Cr(III) carcinogenicity;
- In small amounts, Cr(III) is an essential micronutrient for lipid, protein and fat metabolism in animals and humans.


# Traditional soil remediation techniques

- Soil flushing
- Soil washing
- Solidification/Stabilization
- Excavation and landfilling
- Vitrification
- Incineration
- Electrokinetic treatment

*In most cases, these techniques are expensive and technically limited to highly polluted and relatively small areas. Therefore, when vast areas of land have been contaminated, alternative remediation strategies are necessary.*

# Soil phytoremediation

The use of plants for the in situ remediation of soil contaminated with organic or inorganic pollutants



# Phytoextraction

1. Extraction and accumulation of metals in plant tissues;
2. Harvesting of the above ground plant material;
3. Recovery of high-price metals from the harvested plants (phytomining), or  
Burning of the harvested plants and disposal under controlled conditions (usual metals).

# Basic conditions of plants used for metal phytoextraction

- Tolerate, uptake and accumulate high metal concentrations (hyperaccumulator plants);
- High soil-to-shoot transfer factors;
- High biomass production;
- Fast grow.

# Hyperaccumulator plants

Disadvantages:

- High percentage of hyperaccumulators are found exclusively in tropical regions; therefore, they don't grow, or grow slowly, under temperate climate;
- Low biomass;
- From over 400 hyperaccumulator species identified, more than 80% accumulate Ni rather than other metals.

*Most hyperaccumulator plants cannot remove large quantities of heavy metals in a given period of time per unit of land area.*

*Species that accumulate lower metal concentrations but are high biomass producers may also be used for phytoextraction.*

# Cr(VI) hyperaccumulator plants

- *Leptospermum scoparium* (New Zealand tea tree) (Lyon et al., 1971);
- *Commelina communis* L (Asiatic dayflower) (Tang et al., 2002);
- *Pteris vittata* (Chinese brake fern) (Su et al., 2005);
- *Salsola kali* (Tumbleweed) (Gardea-Torresdey et al., 2005);
- *Leersia hexandra* Swartz (Southern cutgrass) (Zhang et al., 2007);
- *Brassica juncea* (Indian mustard) (Hema et al., 2008).

# Objective of the study



To investigate the capacity of *Zea mays* (corn), a crop plant with large yields and fast growth, to remediate hexavalent chromium polluted soils.

# Materials and methods

- Soil sample was collected from the arable horizon (0-30 cm depth) of a public garden located in Timisoara;
- The experiment was conducted in 500 mL plastic pots filled with 300 g of homogenized air-dried soil;
- Potassium dichromate solution was applied to soil, yielding the concentrations of 20, 40 and 60 mg Cr(VI) /kg dw soil;
- The soil was then allowed to equilibrate for a period of 15 days; Afterwards, 10 seeds of *Zea mays* were sown on the surface of each pot;
- Pots were irrigated with water once a week with deionised water;
- After 40 days of Cr(VI) exposure, plants were harvested, separated in different parts (roots, stems and leaves) and cleaned;
- The plant parts were dried in an oven at 80° C for two days and then ashed in a muffle furnace at 600°C for 6 h;
- The ash was dissolved in a mixture of 2 M HCl and 1 M HNO<sub>3</sub> and analyzed for Cr(VI) by the 1,5-diphenylcarbazide colorimetric method.

# Results

Cr(VI) concentration in plant tissues, as a function of Cr(VI) concentration in soil




Figure 1. Cr(VI) concentration in root tissues, as a function of Cr(VI) concentration in soil




Figure 2. Cr(VI) concentration in stem tissues, as a function of Cr(VI) concentration in soil




Figure 3. Cr(VI) concentration in leaf tissues, as a function of Cr(VI) concentration in soil

# Results

Cr(VI) concentration in plant tissues, as a function of plant part




Figure 4. Cr(VI) concentration in plant tissues, at 20 mg Cr(VI)/kg dw soil




Figure 5. Cr(VI) concentration in plant tissues, at 40 mg Cr(VI)/kg dw soil




Figure 6. Cr(VI) concentration in plant tissues, at 60 mg Cr(VI)/kg dw soil

# Conclusions

- Hexavalent chromium is a toxic metal and needs to be removed from polluted soils;
- Most currently available soil remediation technologies are uneconomical; therefore, they are limited to small areas;
- Phytoextraction can represent a cheaper alternative soil remediation method;
- There are no well established Cr(VI) hyperaccumulator plants;
- This study investigated the capacity of *Zea mays* (corn), a crop plant with large yields and fast growth, to remediate hexavalent chromium polluted soils;
- Cr(VI) concentration in plant organs increased with the increase of Cr(VI) soil concentration;
- Cr(VI) was immobilized mainly in roots, and its concentration in plant organs decreased in the following order: roots > stems > leaves;
- Cr(VI) was poor translocated from the roots to the shoot; however, the increase in Cr(VI) soil concentration improved the translocation of Cr(VI) to the above ground plant parts;
- Although *Zea mays* seem to be a low hexavalent chromium-accumulating plant, it could be a potential phytoremediator of contaminated soils because it's a plant specie with large biomass and fast growth.

# Future work

- Since no visible phytotoxic symptoms were observed on *Zea mays* over the Cr(VI) concentration range 20 - 60 mg/kg d.w. soil, separate laboratory studies will be conducted on this plant with the aim to investigate the phytoremediation of soils affected by higher amounts of hexavalent chromium.
- Long term field studies will be conducted, which will allow plants to reach their maturity, in order to determine the capacity of mature plants to extract and accumulate hexavalent chromium from polluted soils.

# References

1. J. Barceló, C. Poschenrieder, Phytoremediation: principles and perspectives, Contributions to science, 2(3), 2003, 333-344.
2. N. Rajakaruna, K.M. Tompkins, P.G. Pavicevic, Phytoremediation: an affordable green technology for the clean-up of metal-contaminated sites in Sri Lanka, Cey. J. Sci. (Bio. Sci.) 35 (1), 2006, 25-39.
3. E. Pilon-Smits, Phytoremediation, Annu. Rev. Plant. Biol., 56, 2005, 15-39.
4. L. R. Hossner, R. H. Loeppert, R. J. Newton, P. J. Szaniszlo, M. Attrep, Jr., Phytoaccumulation of chromium, uranium, and plutonium in plant systems, Amarillo National Resource Center for Plutonium, 1998.
5. Y. Su, F. X. Han, B.B. M. Sridhar, D. L. Monts, Phytotoxicity and phytoaccumulation of trivalent and hexavalent chromium in brake fern, Environmental Toxicology and Chemistry, 24 (8), 2005, 2019-2026.
6. G.L. Lyon, P.J. Peterson, R.R. Brooks, G.W. Butler. Calcium, magnesium and trace elements in a New Zealand serpentine flora. J. Ecol., 59, 1971, 421-429.
7. X.-H. Zhang, J. Liu, H.-T. Huang, J. Chen, Y.-N. Zhu, D.-Q. Wang, Chromium accumulation by the hyperaccumulator plant *Leersia hexandra* Swartz, Chemosphere, 67 (6), 2007, 1138-1143.
8. D. Hema, A. Altaf, I. Muhammad, Genotypic variation in the phytoremediation potential of Indian Mustard for chromium, Environmental management, 41 (5), 2008, 734-741.
9. Y. Su, F.X. Han, B.B.M. Sridhar, D.L. Monts, Phytotoxicity and phytoaccumulation of trivalent and hexavalent chromium in brake fern, Environmental Toxicology and Chemistry, 24 (8), 2005, 2019-2026.
10. J.L. Gardea-Torresdey, G. de la Rosa, J.R. Peralta-Videa, M. Montes, G. Cruz-Jimenez, I. Cano-Aguilera, Differential uptake and transport of trivalent and hexavalent chromium by tumbleweed (*Salsola kali*). Arch. Environ. Contam. Toxicol, 48 (2), 2005, 225-232
11. S. Tang, L. Xi, Accumulation of chromium by *Commelina communis* L. grown in solution with different concentrations of Cr and L-histidine, Journal of Zhejiang University SCIENCE, 3 (2), 2002, 232 -236.
12. S. Pantawat, P. Wasant, K. Sutha, K. Eakalak, Chromium Removal from Soil by Phytoremediation with Weed Plant Species in Thailand, Water Air and Soil Pollution, 6 (1-2), 2006, 191-206
13. J.L. Gardea-Torresdey, M.V. Aldrich, J.R. Peralta-Videa, J.G. Parsons, Reduction of health risks due to chromium(VI) using Mesquite: a potential Cr phytoremediator, Proceedings of the 227th American Chemical Society Meeting, 28 March - 1 April 2004, Anaheim, California. p 461-466, 2004.
14. A.C. Zayed, N. Terry, Chromium in the environment: Factors affecting biological remediation, Plant Soil, 249, 2003, 139-156.